

Lampiran 7. Hasil Uji Analisis Data Karakteristik Responden

Uji Chi-square

1a. Gravida * Frekuensi ANC

Crosstab

			Frekuensi		Total
			2 - 3	>= 4	
Gravida	Primi	Count	1	19	20
		% within Gravida	5.0%	95.0%	100.0%
	Sekundi	Count	1	18	19
		% within Gravida	5.3%	94.7%	100.0%
	Multi	Count	0	7	7
		% within Gravida	.0%	100.0%	100.0%
Total	Count	2	44	46	
	% within Gravida	4.3%	95.7%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.377 ^a	2	.828
Likelihood Ratio	.678	2	.713
Linear-by-Linear Association	.191	1	.662
N of Valid Cases	46		

a. 3 cells (50.0%) have expected count less than 5. The minimum expected count is .30.

1b. Usia Saat Hamil * Frekuensi ANC

Crosstab

			Frekuensi		Total
			2 - 3	>= 4	
Usia SH	20 - 25	Count	1	20	21
		% within Usia SH	4.8%	95.2%	100.0%
	26 - 30	Count	0	11	11
		% within Usia SH	.0%	100.0%	100.0%
	31 - 35	Count	1	13	14
		% within Usia SH	7.1%	92.9%	100.0%
Total	Count	2	44	46	
	% within Usia SH	4.3%	95.7%	100.0%	



Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.772 ^a	2	.680
Likelihood Ratio	1.208	2	.547
Linear-by-Linear Association	.064	1	.800
N of Valid Cases	46		

a. 3 cells (50.0%) have expected count less than 5. The minimum expected count is .48.

1c. Usia Menikah * Frekuensi ANC

Crosstab

			Frekuensi		Total
			2 - 3	>= 4	
Usia M	< 20	Count	0	5	5
		% within Usia M	.0%	100.0%	100.0%
	>= 20	Count	2	39	41
		% within Usia M	4.9%	95.1%	100.0%
Total		Count	2	44	46
		% within Usia M	4.3%	95.7%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.255 ^b	1	.614		
Continuity Correction ^a	.000	1	1.000		
Likelihood Ratio	.471	1	.492		
Fisher's Exact Test				1.000	.792
Linear-by-Linear Association	.249	1	.617		
N of Valid Cases	46				

a. Computed only for a 2x2 table

b. 3 cells (75.0%) have expected count less than 5. The minimum expected count is .22.



1d. Sumber Informasi * Frekuensi ANC

Crosstab

			Frekuensi		Total
			2 - 3	>= 4	
Sumber	Dokter	Count	0	7	7
		% within Sumber	.0%	100.0%	100.0%
	Bidan	Count	2	35	37
		% within Sumber	5.4%	94.6%	100.0%
	Ahli Gizi	Count	0	2	2
		% within Sumber	.0%	100.0%	100.0%
Total		Count	2	44	46
		% within Sumber	4.3%	95.7%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.509 ^a	2	.775
Likelihood Ratio	.893	2	.640
Linear-by-Linear Association	.131	1	.717
N of Valid Cases	46		

a. 4 cells (66.7%) have expected count less than 5. The minimum expected count is .09.

2a. Pendidikan * Pengetahuan Anemia

Crosstab

			Pengetahuan			Total
			Kurang	Cukup	Baik	
Pendidikan	Rendah	Count	5	12	0	17
		% within Pendidikan	29.4%	70.6%	.0%	100.0%
	Menengah	Count	10	14	2	26
		% within Pendidikan	38.5%	53.8%	7.7%	100.0%
	Tinggi	Count	0	3	0	3
		% within Pendidikan	.0%	100.0%	.0%	100.0%
Total		Count	15	29	2	46
		% within Pendidikan	32.6%	63.0%	4.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.995 ^a	4	.407
Likelihood Ratio	5.618	4	.230
Linear-by-Linear Association	.234	1	.629
N of Valid Cases	46		

a. 5 cells (55.6%) have expected count less than 5. The minimum expected count is .13.

2b. Usia Saat Hamil * Pengetahuan Anemia

Crosstab

			Pengetahuan			Total
			Kurang	Cukup	Baik	
Usia SH	20 - 25	Count	9	12	0	21
		% within Usia SH	42.9%	57.1%	.0%	100.0%
	26 - 30	Count	2	8	1	11
		% within Usia SH	18.2%	72.7%	9.1%	100.0%
	31 - 35	Count	4	9	1	14
		% within Usia SH	28.6%	64.3%	7.1%	100.0%
Total	Count	15	29	2	46	
	% within Usia SH	32.6%	63.0%	4.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.465 ^a	4	.483
Likelihood Ratio	4.272	4	.370
Linear-by-Linear Association	1.610	1	.205
N of Valid Cases	46		

a. 5 cells (55.6%) have expected count less than 5. The minimum expected count is .48.

2c. Usia Menikah * Pengetahuan Anemia

Crosstab

			Pengetahuan			Total
			Kurang	Cukup	Baik	
Usia M	< 20	Count	1	4	0	5
		% within Usia M	20.0%	80.0%	.0%	100.0%
	>= 20	Count	14	25	2	41
		% within Usia M	34.1%	61.0%	4.9%	100.0%
Total	Count	15	29	2	46	
	% within Usia M	32.6%	63.0%	4.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.773 ^a	2	.679
Likelihood Ratio	1.011	2	.603
Linear-by-Linear Association	.129	1	.719
N of Valid Cases	46		

a. 4 cells (66.7%) have expected count less than 5. The minimum expected count is .22.

2d. Sumber Informasi * Pengetahuan Anemia

Crosstab

			Pengetahuan			Total
			Kurang	Cukup	Baik	
Sumber	Dokter	Count	3	2	2	7
		% within Sumber	42.9%	28.6%	28.6%	100.0%
	Bidan	Count	12	25	0	37
		% within Sumber	32.4%	67.6%	.0%	100.0%
	Ahli Gizi	Count	0	2	0	2
		% within Sumber	.0%	100.0%	.0%	100.0%
Total	Count	15	29	2	46	
	% within Sumber	32.6%	63.0%	4.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	13.894 ^a	4	.008
Likelihood Ratio	11.185	4	.025
Linear-by-Linear Association	.068	1	.794
N of Valid Cases	46		

a. 7 cells (77.8%) have expected count less than 5. The minimum expected count is .09.

2e. Tempat ANC * Pengetahuan Anemia

Crosstab

			Pengetahuan			Total
			Kurang	Cukup	Baik	
ANC	Prak Dokter	Count	2	0	2	4
		% within ANC	50.0%	.0%	50.0%	100.0%
	Prak Bidan	Count	9	17	0	26
		% within ANC	34.6%	65.4%	.0%	100.0%
	RS	Count	1	2	0	3
		% within ANC	33.3%	66.7%	.0%	100.0%
	Puskesmas	Count	3	10	0	13
		% within ANC	23.1%	76.9%	.0%	100.0%
	Total	Count	15	29	2	46
		% within ANC	32.6%	63.0%	4.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	24.714 ^a	6	.000
Likelihood Ratio	15.966	6	.014
Linear-by-Linear Association	.000	1	.986
N of Valid Cases	46		

a. 9 cells (75.0%) have expected count less than 5. The minimum expected count is .13.