

Lampiran 5. SPSS

Karakteristik Usia dengan Osteoartritis Lutut

keluhan * usia Crosstabulation

			usia						Total
			15-24	25-34	35-44	45-54	55-64	>64	
keluhan OA	Count		3	14	29	81	80	89	296
	% within keluhan		1,0%	4,7%	9,8%	27,4%	27,0%	30,1%	100,0%
	% within usia		1,9%	4,6%	6,4%	17,0%	20,2%	32,2%	14,3%
	% of Total		0,1%	0,7%	1,4%	3,9%	3,9%	4,3%	14,3%
non OA	Count		159	291	422	396	316	187	1771
	% within keluhan		9,0%	16,4%	23,8%	22,4%	17,8%	10,6%	100,0%
	% within usia		98,1%	95,4%	93,6%	83,0%	79,8%	67,8%	85,7%
	% of Total		7,7%	14,1%	20,4%	19,2%	15,3%	9,0%	85,7%
Total	Count		162	305	451	477	396	276	2067
	% within keluhan		7,8%	14,8%	21,8%	23,1%	19,2%	13,4%	100,0%
	% within usia		100,0%	100,0%	100,0%	100,0%	100,0%	100,0%	100,0%
	% of Total		7,8%	14,8%	21,8%	23,1%	19,2%	13,4%	100,0%

Karakteristik Jenis Kelamin dengan Osteoartritis Lutut

keluhan * jeniskelamin Crosstabulation

			jeniskelamin		Total
			laki-laki	perempuan	
keluhan OA	Count		73	223	296
	% within keluhan		24,7%	75,3%	100,0%
	% within jeniskelamin		9,8%	16,8%	14,3%

	% within keluhan	37,8%	62,2%	100,0%
	% within jeniskelamin	90,2%	83,2%	85,7%
	% of Total	32,4%	53,3%	85,7%
Total	Count	742	1325	2067
	% within keluhan	35,9%	64,1%	100,0%
	% within jeniskelamin	100,0%	100,0%	100,0%
	% of Total	35,9%	64,1%	100,0%

Karakteristik Usia dengan Osteoartritis Lutut pada Jenis Kelamin Laki-laki

keluhan * usia Crosstabulation^a

			usia						Total
			15-24	25-34	35-44	45-54	55-64	>64	
keluhan	OA	Count	1	6	7	18	21	20	73
		% within keluhan	1,4%	8,2%	9,6%	24,7%	28,8%	27,4%	100,0%
		% within usia	1,6%	5,8%	4,1%	10,3%	14,9%	21,7%	9,8%
		% of Total	0,1%	0,8%	0,9%	2,4%	2,8%	2,7%	9,8%
non	OA	Count	61	97	162	157	120	72	669
		% within keluhan	9,1%	14,5%	24,2%	23,5%	17,9%	10,8%	100,0%
		% within usia	98,4%	94,2%	95,9%	89,7%	85,1%	78,3%	90,2%
		% of Total	8,2%	13,1%	21,8%	21,2%	16,2%	9,7%	90,2%
Total	Count	62	103	169	175	141	92	742	
	% within keluhan	8,4%	13,9%	22,8%	23,6%	19,0%	12,4%	100,0%	
	% within usia	100,0%	100,0%	100,0%	100,0%	100,0%	100,0%	100,0%	
	% of Total	8,4%	13,9%	22,8%	23,6%	19,0%	12,4%	100,0%	

a. jeniskelamin = laki-laki

Karakteristik Usia dengan Osteoartritis Lutut pada Jenis Kelamin Perempuan

keluhan * usia Crosstabulation^a

			usia						Total
			15-24	25-34	35-44	45-54	55-64	>64	
keluhan	OA	Count	2	8	22	63	59	69	223
		% within keluhan	0,9%	3,6%	9,9%	28,3%	26,5%	30,9%	100,0%
		% within usia	2,0%	4,0%	7,8%	20,9%	23,1%	37,5%	16,8%
		% of Total	0,2%	0,6%	1,7%	4,8%	4,5%	5,2%	16,8%
non OA		Count	98	194	260	239	196	115	1102
		% within keluhan	8,9%	17,6%	23,6%	21,7%	17,8%	10,4%	100,0%
		% within usia	98,0%	96,0%	92,2%	79,1%	76,9%	62,5%	83,2%
		% of Total	7,4%	14,6%	19,6%	18,0%	14,8%	8,7%	83,2%
Total		Count	100	202	282	302	255	184	1325
		% within keluhan	7,5%	15,2%	21,3%	22,8%	19,2%	13,9%	100,0%
		% within usia	100,0%	100,0%	100,0%	100,0%	100,0%	100,0%	100,0%
		% of Total	7,5%	15,2%	21,3%	22,8%	19,2%	13,9%	100,0%

a. jeniskelamin = perempuan

Faktor resiko antara usia dengan osteoartritis lutut pada seluruh subjek

usia * keluhan Crosstabulation

			keluhan		Total
			OA	non OA	
usia	25-34	Count	14	291	305
		% within usia	4,6%	95,4%	100,0%
		% within keluhan	82,4%	64,7%	65,3%
		% of Total	3,0%	62,3%	65,3%
	15-24	Count	3	159	162

	% of Total	0,6%	34,0%	34,7%
Total	Count	17	450	467
	% within usia	3,6%	96,4%	100,0%
	% within keluhan	100,0%	100,0%	100,0%
	% of Total	3,6%	96,4%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2,262 ^a	1	,133		
Continuity Correction ^b	1,548	1	,213		
Likelihood Ratio	2,519	1	,112		
Fisher's Exact Test				,194	,103
Linear-by-Linear Association	2,257	1	,133		
N of Valid Cases	467				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 5,90.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (25-34 / 15-24)	2,550	,722	9,006
For cohort keluhan = OA	2,479	,723	8,500
For cohort keluhan = non OA	,972	,941	1,004
N of Valid Cases	467		

usia * keluhan Crosstabulation

			keluhan		Total
			OA	non OA	
usia	35-44	Count	29	422	451
		% within usia	6,4%	93,6%	100,0%
		% within keluhan	90,6%	72,6%	73,6%
		% of Total	4,7%	68,8%	73,6%
	15-24	Count	3	159	162
		% within usia	1,9%	98,1%	100,0%
		% within keluhan	9,4%	27,4%	26,4%
		% of Total	0,5%	25,9%	26,4%
Total		Count	32	581	613
		% within usia	5,2%	94,8%	100,0%
		% within keluhan	100,0%	100,0%	100,0%
		% of Total	5,2%	94,8%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	5,049 ^a	1	,025		
Continuity Correction ^b	4,166	1	,041		
Likelihood Ratio	6,134	1	,013		
Fisher's Exact Test				,023	,015
Linear-by-Linear Association	5,041	1	,025		
N of Valid Cases	613				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 8,46.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (35-44 / 15-24)	3,642	1,094	12,124
For cohort keluhan = OA	3,472	1,072	11,244
For cohort keluhan = non OA	,953	,923	,984
N of Valid Cases	613		

usia * keluhan Crosstabulation

			keluhan		Total
			OA	non OA	
usia 45-54	Count	81	396	477	
	% within usia	17,0%	83,0%	100,0%	
	% within keluhan	96,4%	71,4%	74,6%	
	% of Total	12,7%	62,0%	74,6%	
15-24	Count	3	159	162	
	% within usia	1,9%	98,1%	100,0%	
	% within keluhan	3,6%	28,6%	25,4%	
	% of Total	0,5%	24,9%	25,4%	
Total	Count	84	555	639	
	% within usia	13,1%	86,9%	100,0%	
	% within keluhan	100,0%	100,0%	100,0%	
	% of Total	13,1%	86,9%	100,0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	24,244 ^a	1	,000		
Continuity Correction ^b	22,937	1	,000		
Likelihood Ratio	32,818	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	24,206	1	,000		
N of Valid Cases	639				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 21,30.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (45-54 / 15-24)	10,841	3,375	34,823
For cohort keluhan = OA	9,170	2,937	28,628
For cohort keluhan = non OA	,846	,808	,885
N of Valid Cases	639		

usia * keluhan Crosstabulation

			keluhan		Total
			OA	non OA	
usia	55-64	Count	80	316	396
		% within usia	20,2%	79,8%	100,0%
		% within keluhan	96,4%	66,5%	71,0%
		% of Total	14,3%	56,6%	71,0%

	% within keluhan	3,6%	33,5%	29,0%
	% of Total	0,5%	28,5%	29,0%
Total	Count	83	475	558
	% within usia	14,9%	85,1%	100,0%
	% within keluhan	100,0%	100,0%	100,0%
	% of Total	14,9%	85,1%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	30,574 ^a	1	,000		
Continuity Correction ^b	29,142	1	,000		
Likelihood Ratio	40,903	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	30,519	1	,000		
N of Valid Cases	558				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 24,10.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (55-64 / 15-24)	13,418	4,172	43,156
For cohort keluhan = OA	10,909	3,496	34,043
For cohort keluhan = non OA	,813	,770	,858

usia * keluhan Crosstabulation

			keluhan		Total
			OA	non OA	
usia >64	Count	89	187	276	
	% within usia	32,2%	67,8%	100,0%	
	% within keluhan	96,7%	54,0%	63,0%	
	% of Total	20,3%	42,7%	63,0%	
15-24	Count	3	159	162	
	% within usia	1,9%	98,1%	100,0%	
	% within keluhan	3,3%	46,0%	37,0%	
	% of Total	0,7%	36,3%	37,0%	
Total	Count	92	346	438	
	% within usia	21,0%	79,0%	100,0%	
	% within keluhan	100,0%	100,0%	100,0%	
	% of Total	21,0%	79,0%	100,0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	56,836 ^a	1	,000		
Continuity Correction ^b	55,019	1	,000		
Likelihood Ratio	73,352	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	56,706	1	,000		
N of Valid Cases	438				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 34,03.

b. Computed only for a 2x2 table

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (>64 / 15-24)	25,225	7,831	81,254
For cohort keluhan = OA	17,413	5,602	54,122
For cohort keluhan = non OA	,690	,635	,751
N of Valid Cases	438		

Faktor Resiko antara Usia dengan Osteoartritis Lutut pada Jenis Kelamin Laki-laki dan Perempuan

usia * keluhan Crosstabulation^a

			keluhan		Total
			OA	non OA	
usia	25-34	Count	6	97	103
		% within usia	5,8%	94,2%	100,0%
		% within keluhan	85,7%	61,4%	62,4%
		% of Total	3,6%	58,8%	62,4%
	15-24	Count	1	61	62
		% within usia	1,6%	98,4%	100,0%
		% within keluhan	14,3%	38,6%	37,6%
		% of Total	0,6%	37,0%	37,6%
Total		Count	7	158	165
		% within usia	4,2%	95,8%	100,0%
		% within keluhan	100,0%	100,0%	100,0%
		% of Total	4,2%	95,8%	100,0%

a. jeniskelamin = laki-laki

Chi-Square Tests^a

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1,690 ^b	1	,194	,257	,187
Continuity Correction ^c	,813	1	,367		
Likelihood Ratio	1,942	1	,163		
Fisher's Exact Test					
Linear-by-Linear Association	1,680	1	,195		
N of Valid Cases	165				

a. jeniskelamin = laki-laki

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,63.

c. Computed only for a 2x2 table

Risk Estimate^a

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (25-34 / 15-24)	3,773	,443	32,105
For cohort keluhan = OA	3,612	,445	29,300
For cohort keluhan = non OA	,957	,904	1,014
N of Valid Cases	165		

a. jeniskelamin = laki-laki

usia * keluhan Crosstabulation^a

			keluhan		Total
			OA	non OA	
usia	25-34	Count	8	194	202
		% within usia	4,0%	96,0%	100,0%

15-24	Count	2	98	100
	% within usia	2,0%	98,0%	100,0%
	% within keluhan	20,0%	33,6%	33,1%
	% of Total	0,7%	32,5%	33,1%
Total	Count	10	292	302
	% within usia	3,3%	96,7%	100,0%
	% within keluhan	100,0%	100,0%	100,0%
	% of Total	3,3%	96,7%	100,0%

a. jeniskelamin = perempuan

Chi-Square Tests^a

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	,803 ^b	1	,370		
Continuity Correction ^c	,307	1	,579		
Likelihood Ratio	,874	1	,350		
Fisher's Exact Test				,506	,300
Linear-by-Linear Association	,800	1	,371		
N of Valid Cases	302				

a. jeniskelamin = perempuan

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 3,31.

c. Computed only for a 2x2 table

Risk Estimate^a

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (25-34 / 15-24)	2,021	,421	9,697
For cohort keluhan = OA	1,980	,428	9,153
For cohort keluhan = non OA	,980	,942	1,020
N of Valid Cases	302		

a. jeniskelamin = perempuan

usia * keluhan Crosstabulation^a

			keluhan		Total
			OA	non OA	
usia	35-44	Count	7	162	169
		% within usia	4,1%	95,9%	100,0%
		% within keluhan	87,5%	72,6%	73,2%
		% of Total	3,0%	70,1%	73,2%
	15-24	Count	1	61	62
		% within usia	1,6%	98,4%	100,0%
		% within keluhan	12,5%	27,4%	26,8%
		% of Total	0,4%	26,4%	26,8%
Total		Count	8	223	231
		% within usia	3,5%	96,5%	100,0%
		% within keluhan	100,0%	100,0%	100,0%
		% of Total	3,5%	96,5%	100,0%

a. jeniskelamin = laki-laki

Chi-Square Tests^a

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,868 ^b	1	,352		
Continuity Correction ^c	,276	1	,599		
Likelihood Ratio	1,007	1	,316		
Fisher's Exact Test				,686	,318
Linear-by-Linear Association	,864	1	,353		
N of Valid Cases	231				

a. jeniskelamin = laki-laki

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,15.

c. Computed only for a 2x2 table

Risk Estimate^a

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (35-44 / 15-24)	2,636	,318	21,869
For cohort keluhan = OA	2,568	,322	20,453
For cohort keluhan = non OA	,974	,932	1,019
N of Valid Cases	231		

a. jeinskelamin = laki-laki

usia * keluhan Crosstabulation^a

			keluhan		Total
			OA	non OA	
usia	35-44	Count	22	260	282
		% within usia	7,8%	92,2%	100,0%

15-24	Count	2	98	100
	% within usia	2,0%	98,0%	100,0%
	% within keluhan	8,3%	27,4%	26,2%
	% of Total	0,5%	25,7%	26,2%
Total	Count	24	358	382
	% within usia	6,3%	93,7%	100,0%
	% within keluhan	100,0%	100,0%	100,0%
	% of Total	6,3%	93,7%	100,0%

a. jeniskelamin = perempuan

Chi-Square Tests^a

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	4,220 ^b	1	,040		
Continuity Correction ^c	3,292	1	,070		
Likelihood Ratio	5,210	1	,022		
Fisher's Exact Test				,052	,027
Linear-by-Linear Association	4,209	1	,040		
N of Valid Cases	382				

a. jeniskelamin = perempuan

b. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 6,28.

c. Computed only for a 2x2 table

Risk Estimate^a

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (35-44 / 15-24)	4,146	,957	17,962
For cohort keluhan = OA	3,901	,934	16,291
For cohort keluhan = non OA	,941	,900	,983
N of Valid Cases	382		

a. jeniskelamin = perempuan

usia * keluhan Crosstabulation^a

			keluhan		Total
			OA	non OA	
usia	45-54	Count	18	157	175
		% within usia	10,3%	89,7%	100,0%
		% within keluhan	94,7%	72,0%	73,8%
		% of Total	7,6%	66,2%	73,8%
	15-24	Count	1	61	62
		% within usia	1,6%	98,4%	100,0%
		% within keluhan	5,3%	28,0%	26,2%
		% of Total	0,4%	25,7%	26,2%
Total		Count	19	218	237
		% within usia	8,0%	92,0%	100,0%
		% within keluhan	100,0%	100,0%	100,0%
		% of Total	8,0%	92,0%	100,0%

a. jeniskelamin = laki-laki

Chi-Square Tests^a

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4,670 ^b	1	,031		
Continuity Correction ^c	3,568	1	,059		
Likelihood Ratio	6,133	1	,013		
Fisher's Exact Test				,030	,020
Linear-by-Linear Association	4,650	1	,031		
N of Valid Cases	237				

a. jeniskelamin = laki-laki

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,97.

c. Computed only for a 2x2 table

Risk Estimate^a

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (45-54 / 15-24)	6,994	,914	53,531
For cohort keluhan = OA	6,377	,869	46,780
For cohort keluhan = non OA	,912	,859	,968
N of Valid Cases	237		

a. jeniskelamin = laki-laki

usia * keluhan Crosstabulation^a

			keluhan		Total
			OA	non OA	
usia	45-54	Count	63	239	302
		% within usia	20,9%	79,1%	100,0%

15-24	Count	2	98	100
	% within usia	2,0%	98,0%	100,0%
	% within keluhan	3,1%	29,1%	24,9%
	% of Total	0,5%	24,4%	24,9%
Total	Count	65	337	402
	% within usia	16,2%	83,8%	100,0%
	% within keluhan	100,0%	100,0%	100,0%
	% of Total	16,2%	83,8%	100,0%

a. jeniskelamin = perempuan

Chi-Square Tests^a

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	19,716 ^b	1	,000		
Continuity Correction ^c	18,349	1	,000		
Likelihood Ratio	26,820	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	19,667	1	,000		
N of Valid Cases	402				

a. jeniskelamin = perempuan

b. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 16,17.

c. Computed only for a 2x2 table

Risk Estimate^a

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (45-54 / 15-24)	12,916	3,100	53,824
For cohort keluhan = OA	10,430	2,599	41,854
For cohort keluhan = non OA	,808	,757	,861
N of Valid Cases	402		

a. jeniskelamin = perempuan

usia * keluhan Crosstabulation^a

			keluhan		Total
			OA	non OA	
usia	55-64	Count	21	120	141
		% within usia	14,9%	85,1%	100,0%
		% within keluhan	95,5%	66,3%	69,5%
		% of Total	10,3%	59,1%	69,5%
	15-24	Count	1	61	62
		% within usia	1,6%	98,4%	100,0%
		% within keluhan	4,5%	33,7%	30,5%
		% of Total	0,5%	30,0%	30,5%
Total		Count	22	181	203
		% within usia	10,8%	89,2%	100,0%
		% within keluhan	100,0%	100,0%	100,0%
		% of Total	10,8%	89,2%	100,0%

a. jeniskelamin = laki-laki

Chi-Square Tests^a

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	7,860 ^b	1	,005		
Continuity Correction ^c	6,546	1	,011		
Likelihood Ratio	10,379	1	,001		
Fisher's Exact Test				,003	,002
Linear-by-Linear Association	7,822	1	,005		
N of Valid Cases	203				

a. jeniskelamin = laki-laki

b. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 6,72.

c. Computed only for a 2x2 table

Risk Estimate^a

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (55-64 / 15-24)	10,675	1,403	81,250
For cohort keluhan = OA	9,234	1,270	67,130
For cohort keluhan = non OA	,865	,802	,933
N of Valid Cases	203		

a. jeniskelamin = laki-laki

usia * keluhan Crosstabulation^a

			keluhan		Total
			OA	non OA	
usia	55-64	Count	59	196	255
		% within usia	23,1%	76,9%	100,0%
		% within keluhan	96,7%	66,7%	71,8%
		% of Total	16,6%	55,3%	71,9%

	% within usia	2,0%	98,0%	100,0%
	% within keluhan	3,3%	33,3%	28,2%
	% of Total	0,6%	27,6%	28,2%
Total	Count	61	294	355
	% within usia	17,2%	82,8%	100,0%
	% within keluhan	100,0%	100,0%	100,0%
	% of Total	17,2%	82,8%	100,0%

a. jeniskelamin = perempuan

Chi-Square Tests^a

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	22,552 ^b	1	,000		
Continuity Correction ^c	21,091	1	,000		
Likelihood Ratio	30,250	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	22,489	1	,000		
N of Valid Cases	355				

a. jeniskelamin = perempuan

b. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 17,18.

c. Computed only for a 2x2 table

Risk Estimate^a

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (55-64 / 15-24)	14,750	3,530	61,630
For cohort keluhan = OA	11,569	2,881	46,451
For cohort keluhan = non OA	,784	,729	,844
N of Valid Cases	355		

Usia * keluhan Crosstabulation^a

			keluhan		Total
			OA	non OA	
usia >64	Count	20	72	92	
	% within usia	21,7%	78,3%	100,0%	
	% within keluhan	95,2%	54,1%	59,7%	
	% of Total	13,0%	46,8%	59,7%	
15-24	Count	1	61	62	
	% within usia	1,6%	98,4%	100,0%	
	% within keluhan	4,8%	45,9%	40,3%	
	% of Total	0,6%	39,6%	40,3%	
Total	Count	21	133	154	
	% within usia	13,6%	86,4%	100,0%	
	% within keluhan	100,0%	100,0%	100,0%	
	% of Total	13,6%	86,4%	100,0%	

a. jeniskelamin = laki-laki

Chi-Square Tests^a

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	12,740 ^b	1	,000		
Continuity Correction ^c	11,088	1	,001		
Likelihood Ratio	16,101	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	12,657	1	,000		
N of Valid Cases	154				

a. jeniskelamin = laki-laki

b. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 8,45.

c. Computed only for a 2x2 table

Risk Estimate^a

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (>64 / 15-24)	16,944	2,210	129,938
For cohort keluhan = OA	13,478	1,857	97,852
For cohort keluhan = non OA	,795	,711	,890
N of Valid Cases	154		

a. jeniskelamin = laki-laki

usia * keluhan Crosstabulation^a

			keluhan		Total
			OA	non OA	
usia >64	Count	69	115	184	
	% within usia	37,5%	62,5%	100,0%	
	% within keluhan	97,2%	54,0%	64,8%	
	% of Total	24,3%	40,5%	64,8%	
15-24	Count	2	98	100	
	% within usia	2,0%	98,0%	100,0%	
	% within keluhan	2,8%	46,0%	35,2%	
	% of Total	0,7%	34,5%	35,2%	
Total	Count	71	213	284	
	% within usia	25,0%	75,0%	100,0%	
	% within keluhan	100,0%	100,0%	100,0%	
	% of Total	25,0%	75,0%	100,0%	

a. jeniskelamin = perempuan

Chi-Square Tests^a

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	43,547 ^b	1	,000		
Continuity Correction ^c	41,674	1	,000		
Likelihood Ratio	56,343	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	43,393	1	,000		
N of Valid Cases	284				

a. jeniskelamin = perempuan

b. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 25,00.

c. Computed only for a 2x2 table

Risk Estimate^a

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (>64 / 15-24)	29,400	7,026	123,030
For cohort keluhan = OA	18,750	4,695	74,873
For cohort keluhan = non OA	,638	,568	,716
N of Valid Cases	284		

a. jeniskelamin = perempuan