

## DAFTAR PUSTAKA

- Brookes G.B. 1985. Vitamin D deficiency and deafness: 1984 update. *Otology & Neurotology*, 6, 102-107.
- Brookes G.B. 1983. Vitamin D deficiency—a new cause of cochlear deafness. *The Journal of Laryngology & Otology*, 97, 405-420.
- Bosworth C. and Boer D. 2013. Impaired vitamin D metabolism in CKD. *Seminars in nephrology*. Elsevier.
- Cankaya E., Bilen Y., Keles M., Uyanik A., Akbas, M., Gungo A., Arslan S & Aydinli B. 2015. Comparison of Serum Vitamin D Levels Among Patients With Chronic Kidney Disease, Patients in Dialysis, and Renal Transplant Patients. *Transplantation proceedings*. Elsevier.
- Cadha S. 2015. International perspective: reducing hearing loss. *J Prim Health Care*, 7, 4.
- Cuna, Battagliano V., Capeli P., SALA I., Donati E., Cianciol G., & Mannad L.A., 2015. Hypoacusia and chronic renal dysfunction: new etiopathogenetic prospective. *Ther Apher Dial*, 19, 111-8.
- Dinghra P. 2013. *Diseases of Ear, Nose and Throat*, 6/e, Elsevier India.
- Dusso A.S. 2011. Kidney disease and vitamin D levels: 25-hydroxyvitamin D, 1,25-dihydroxyvitamin D, and VDR activation. *Kidney International Supplements*, 1, 136-141.
- Dusso A., Gonzales E.A. & MARTIN K.J. 2011. Vitamin D in chronic kidney disease. *Best practice & research Clinical endocrinology & metabolism*, 25, 647-655.
- Erkoch M.F., BULUT S., Imamoglu H., Gumus C. & Kayatas M. 2013. CT assessment of bone remodeling in the otic capsule in chronic renal failure: association with hearing loss. *American Journal of Roentgenology*, 200, 396-399.
- Flint P.W., Haughey B.H., Lund V.J., Niparko J.K., Richarson M.A., Robbins K.T., Thomas J.R. & Lesperance M.M. 2014. *Cummings Otolaryngology: Head and Neck Surgery, 3-Volume Set*, Elsevier Science Health Science.
- Govender S.M., Govender C.D. & Matthews G. 2013. Cochlear function in patients with chronic kidney disease. *South African Journal of Communication Disorders*, 60, 44-49.

- Holick M.F. (2009) Vitamin D status: measurement, interpretation, and clinical application. *Annals of epidemiology*, 19, 73-78.
- Ikeda K., Kobayashi T., Itoh Z., Kusakari J. & Takasaka T. (1989) Evaluation of vitamin D metabolism in patients with bilateral sensorineural hearing loss. *Am J Otol*, 10, 11-13.
- Ikeda K., Kusakari J., Kobayashi T. & Saito Y. 1987. The effect of vitamin D deficiency on the cochlear potentials and the perilymphatic ionized calcium concentration of rats. *Acta Oto-Laryngologica*, 104, 64-72.
- Jamaldeen J., Basheer A., Sarma A.C. & Kadasamy R. (2015) Prevalence and patterns of hearing loss among chronic kidney disease patients undergoing haemodialysis. *Australas Med J*, 8, 41-6.
- JIN, D-C, YUN, SR, LEE, SW, HAN, SW, KIM, W, PARK, J & KIM, YK (2015) Lessons from 30 years' data of Korean end-stage renal disease registry, 1985–2015. *Kidney research and clinical practice*, 34, 132-139.
- Kang S.H., Jung D.A., CHO K.H., Park J.W., Yoon K.W. & DO J.Y. 2015. The association between metabolic syndrome or chronic kidney disease and hearing thresholds in Koreans: the Korean National Health and Nutrition Examination Survey 2009-2012. *PLoS One*, 10, e0120372.
- Kasper D., Fauci A., Hauser S., Longo D., Jameson J. & Localizo J. 2015. *Harrison's Principles of Internal Medicine 19/E (Vol.1 & Vol.2)*, McGraw-Hill Education.
- Kdoqi & Langman C.B. (2005) K-DOQI Clinical Practice Guidelines for Bone Metabolism and Disease in Chronic Kidney Disease. Saunders.
- Kim J.Y., Lee S.B/, Lee C.H. & Kim H.M. 2015. Hearing loss in postmenopausal women with low bone mineral density. *Auris Nasus Larynx*.
- LEVEY A.S., Coresh J., Balk E., Kausz A.T., Levin A., Steffes M.W., Hogg R.J., Perrone R.D., Lau J. & Eknoyan G. (2003) National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Annals of*
- OLUSANYA, BO, NEUMANN, KJ & SAUNDERS, JE (2014) The global burden of disabling hearing impairment: a call to action. *Bulletin of the World Health Organization*, 92, 367-373.
- Levey A.S., Eckardt K.U., Tsukamoto Y., Levin A., Coresh J., Rossert J., Zeeuw D.D., Hostetter T.H., Lameire N. & Eknoyan G. (2005) Definition and classification of

- chronic kidney disease: a position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney international*, 67, 2089-2398.
- Looi L.M., Ganten D., Mcgrath P.F., Gross M. & Griffin G.E. 2015. Hearing loss: a global health issue. *The Lancet*, 385, 943-944.
- Monsell E.M. 2004. The mechanism of hearing loss in Paget's disease of bone. *The Laryngoscope*, 114, 598-606.
- Nigwekar S.U., Tamez H. & Thadhani R.I. 2014. Vitamin D and chronic kidney disease-mineral bone disease (CKD-MBD). *BoneKEy Rep*, 3.
- Pavlovic D. Katicic D., Gulin T. & Josipovic J. 2015. Vitamin D in the Patients with Chronic Kidney Disease: When, to Whom and in Which Form. *Materia socio-medica*, 27, 122.
- Phelps K., Mathew R., Stote K., Hewson L. & Hallendeck D. 2012. Relationships Among eGFR, vitamin D metabolites and PTH 1-84 in ckd. *Kidney Research and Clinical Practice*, 31, A66.
- Qazi R.A. & Martin K.J. (2012) Vitamin D in kidney disease: pathophysiology and the utility of treatment. *Rheumatic Disease Clinics of North America*, 38, 115-123.
- Rahman R., Akhtar N., Khanam A., Aalam M.R. 2016. Prevalence and Patterns of Hearing Loss among Chronic Kidney Disease of Various Stages in Bangladeshi Patients. *American Journal of Medical Sciences and Medicine*. 2016; 4(3):53-58.
- Seo Y.J., Ko S.B., Ha T.H., Gong T.H., Bong J.P., Park D.J. & Park S.Y. 2015. Association of hearing impairment with chronic kidney disease: a cross-sectional study of the Korean general population. *BMC nephrology*, 16, 154. *internal medicine*, 139, 137-147.
- Thodi C., Thodi E., Danielides V., Pasadakis P. & Vargemeziz V. 2006. Hearing in renal failure. *Nephrology Dialysis Transplantation*, 21, 3023-3030.
- Vilayur E., Gopinath B., Harris D.C., Burlutsky G., McMahon C.M. & Mitchell P. 2010 The association between reduced GFR and hearing loss: a cross-sectional population-based study. *Am J Kidney Dis*, 56, 661-9.
- WHO 2013. Millions of people in the world have hearing loss that can be treated or prevented. *Geneva: WHO*, 1-17.

Zou J., Minasyan A., Keisala T., Zhang Y., Wang J.H., Lou Y.R., Kalueff A., Pyykko I. & Touhima P. 2008. Progressive hearing loss in mice with a mutated vitamin D receptor gene. *Audiology and Neurotology*, 13, 219-230.

